Algebra II / I	Mr. Hansen
2/18/2016	

Name:
-------

# Test on Chapter 5

Graphing calculator and PENCIL are required. If you need more room for rough scratch work, use the final two pages, which will not be graded. If you need more room for work that is to be graded, write OVER and continue on the reverse side. You may use the Rule of 72 once if you wish. Circle all answers.

## Please print your numbers like this:



Nonconforming numbers may be misinterpreted, resulting in loss of points. For most students, it is 4 and 9 that cause trouble. European-style 1 and crossed 7 are also acceptable. A "closed-top pointy 4" is also acceptable. *Note:* All answers, unless otherwise stated, must be correct to at least 3 significant digits.

#### Part I: Logarithmic Detective Work (5 pts. each)

**Instructions:** Given that  $\log 2 \approx 0.3$ , estimate each of the following. No credit (**zero points**) for answers that are not supported by work. Be sure to draw a distinction between the equals sign (=) and the approximately-equals sign ( $\approx$ ). The first one is done for you as an example. You are permitted to use the answers for #1 through #3 as "building blocks" if you wish.

1. log 20

**Solution:**  $\log 20 = \log (2.10) = \log 2 + \log 10 \approx 0.3 + 1 = \boxed{1.3}$ 

- 2. log 5
- $\log 8$
- 4. log 80

#### Part II: Creative Work (8 pts. each)

5. Describe how to estimate 80<sup>60</sup> to the nearest power of 10. The only tool you may use is your calculator. (Do use it.) Complete sentences are required for full credit. Circle your final answer.

6. Prove that the base-8 logarithm of any positive real number is always one-third as large as the base-2 logarithm of the same number. You may assume that the change-of-base formula is valid.

#### Part III: Criminal Forensics (20 pts.)

7. Newton's Law of Cooling:  $T(t) = T_S + (T_0 - T_S) e^{-kt}$ , where  $T_S =$  surrounding temperature,  $T_0 =$  initial temperature of object.

Fred Friendly was feeling feverish at home (in his flat on the fifth floor) on the morning of February 15. His body temperature was  $100^{\circ}$  F. when a bullet from a fraudster found Fred and felled him. A famously frugal man, Fred always kept the furnace in his house set to  $55^{\circ}$  F. As he lay dying, Fred frantically fumbled and wrote a note saying that his body temperature was  $100^{\circ}$  F., but he foolishly forgot to record the time or the name of the fraudster who killed him. When police arrived at 11:00 a.m., Fred's corpse had a temperature of  $85^{\circ}$  F., and an hour later, it had cooled to  $82^{\circ}$  F. Compute the time of death to the nearest hour. Show all work clearly. *Hint:* For full credit, you have to find the value of k. However, if that baffles you, check here  $\square$  and use the value 0.1 for k. (That's not correct, but it's close enough to give you the correct final answer.)

Part IV:	Graphing	(15)	pts.	each)	)
----------	----------	------	------	-------	---

Sketch the graph of the $T(t)$ curve from problem #7. Be sure to label your horizontal axis as $t$ an your vertical axis as temperature (° F.). You may use $t = 0$ to denote either the time of death or t time at which the police arrived—your choice, but be sure to indicate which choice you made. F $k$ , use the value you found in #7. (If you were unable to compute $k$ , use the approximate value 0.1.) Mark coordinates for the points where temperature equals 100, 85, and 82 degrees.				
Explain how to sketch the curve $y = 2 \ln (x + 1) - 3$ . ( <i>Note:</i> No credit for saying, "Plug it into measure calculator and make a neat transcription onto paper." You have to describe the steps or transformations that you would perform.) Write your explanation here:				
Sketch the graph and mark coordinates for the points where $x$ equals $0$ , $e-1$ , and $\frac{1}{e}-1$ .				
Asymptotes, if any, must be labeled with their equation and shown as dotted lines.				

#### Part V: Growth/decay problems (10 pts. each)

10. A \$15,000 investment has a yield of approximately 8%, compounded annually. Compute the time required for the investment to be worth \$30,000. Explain your reasoning adequately.

11. A 10 kg lump of radioactive thessoramide (a fictitious substance) has a decay rate of 4% per year. Compute the half-life, i.e., the time needed for the 10 kg lump to decay to 5 kg. Explain your reasoning adequately.

## Rough Scratch Work (NOT GRADED)

## Rough Scratch Work (NOT GRADED)